



## Cautions After vaccination

### ① General cautions after vaccination

- ◆ Avoid heavy exercising for 2 days.
- ◆ Avoid taking a bath for 1 day.
- ◆ Keep clean your vaccinated part.

### ② If you get abnormal symptoms,

- ◆ There can be some symptoms during immunization.
- ◆ Please refer below to check abnormal symptoms.

#### <1> Light symptoms

- ▶ Pain, redness, tenderness on pressure, edemas can be appeared, most of them are disappeared in 2days.
- ▶ Headache, fever, nauseous, myalgia occur seldomly, most of them are occurred in 6~12hours after vaccination and continue for 1~2days, and disappeared.

☞ Symptoms above are temporary. However, if the symptoms continue more than 2 days, please visit doctor.

#### <2> Severe symptoms

- ▶ Anaphylaxis(alergy shock) can occur very seldomly. Anaphylaxis has dyspnea, hoarseness, hives. It occurs right after or in hours after viccination.
- ▶ There is a report for rare possibility of muscular weakness or abnormal sense of finger and toes after vaccination. Relation between symptoms and vaccination are not certified. Most of these symptoms disappeared in 2months.

☞ Those symptoms can be occurred very rarely, if you get these symptoms, visit doctor immediately.

**Ajou Universtiy Infirmary office**  
**(New Student Union Buildint Room#227) ☎ O31)219-1596**